



FAX Page 1 To: Clark County Public Health 360-397-8080 (Confidential FAX Line)

CONFIDENTIAL SEXUALLY TRANSMITTED DISEASE CASE REPORT

TO: Clark County Public Health: STD Program DATE: FROM: FROM PHONE #

Clark County health care providers and health care facilities should use this form to report Sexually Transmitted Disease (STD). Report STDs within three work days (see WAC 246-101-101/301). All information will be strictly managed confidentially. Your cooperation is both encouraged and appreciated.

PATIENT INFORMATION

Form with fields for Patient Information: LAST NAME, FIRST NAME, MIDDLE INITIAL, ADDRESS, TELEPHONE, REASON FOR EXAM, CITY/TOWN, STATE, ZIP CODE, DATE OF DIAGNOSIS, SEX, DATE OF BIRTH, ETHNICITY, RACE, GENDER OF SEX PARTNERS.

DIAGNOSIS - DISEASE

Form for GONORRHEA (lab confirmed) and SYPHILIS. Includes diagnosis, site, treatment, and date tested fields.

Form for CHLAMYDIA TRACHOMATIS (lab confirmed) and HERPES SIMPLEX. Includes diagnosis, site, treatment, and date tested fields.

Form for provider information: DIAGNOSING CLINICIAN, PERSON COMPLETING FORM, FACILITY NAME, ADDRESS, CITY, STATE, TELEPHONE.

PARTNER MANAGEMENT PLAN (Select method of ensuring partner treatment). Includes instructions and a red arrow pointing to 'INSTRUCTIONS On Page 2'.

PRIVILEGED AND CONFIDENTIAL COMMUNICATIONS: The information contained in this message is privileged, confidential, or otherwise exempt from disclosure and is intended solely for the use of the individual(s) named above.

## PARTNER MANAGEMENT PLAN INSTRUCTIONS

### Gonorrhea or Chlamydial Infection: Partner Treatment

All partners should be treated as if they are infected.

If the provider takes responsibility to ensure partner treatment, the provider should examine and treat all patient's sex partners from the previous 60 days.

If this is **not** possible, patients should be offered medication to give to as many of their sex partners as they are able to contact and/or should be referred to Clark County Public Health for partner notification assistance.

Free medication is available for your patient's partner(s).

To obtain **FREE** medication for your patient's partner(s), call or fax a prescription to one of the pharmacies participating in your area. Preprinted prescription are attached to this case report.

**Note: Only participating pharmacies have** stocks of **FREE Public Health medication** to dispense to patients for their partner(s).

Clark County Public Health may also provide free medication to your patient to give to his or her partner(s).

Clark County Public Health recommends that you refer patients with any one or more of the following risks to the health department for help notifying their partners:

- Patient with 2 or more sex partners in the last 60 days , or
- Patient does not think he/she will have sex again with sex partners from the last 60 days, or
- Patient is unable/unwilling to contact one or more partner(s), or
- Patient is a man who has sex with other men

Although Clark County Public Health requests that you refer patients with these risks to us, we also ask that you make every effort to help patients with these risks assure that their partners are treated, either by seeing the partners yourself or by offering patients free medication to give to their partners.

**Complete the partner management plan** on the Confidential Sexually Transmitted Disease Case Report FAX form to define a partner management plan.

For copies of this case report or questions on how to fill it out, call the Clark County Public Health: **360-397-8082**.

### Other STDs: Partner Treatment

All patients with infectious syphilis, chancroid, LGV or granuloma inguinale are routinely contacted by Clark County Public Health. Patients diagnosed with genital herpes should be advised to notify their sex partners and should be informed that their partners should contact their provider for testing.

## RECOMMENDED REGIMENS FOR ANTIMICROBIALS LISTED ON CASE REPORTS\*

#### Gonorrhea (uncomplicated):

Ceftriaxone.....	125-250 mg IM, single dose,	<b>OR</b>
Cefpodoxime.....	400 mg PO, single dose,	<b>OR</b>
Cefixime.....	400 mg PO, single dose,	<b>OR</b>

#### Alternatives:

Azithromycin.....2g PO as a single dose (additional chlamydia therapy not needed)  
Health care providers should no longer use fluoroquinolones (Levofloxacin or Ciprofloxacin) as first line therapy due to increased prevalence of quinolone-resistant *Neisseria gonorrhoeae* (QRNG).

**Plus**, if Chlamydia infection is NOT ruled out

Azithromycin.....	1g PO as a single dose,	<b>OR</b>
Doxycycline.....	100 mg PO BID for 7 days	

#### Chlamydia Trachomatis (uncomplicated):

Azithromycin.....	1g PO single dose,	<b>OR</b>
Doxycycline.....	100 mg PO BID for 7 days,	<b>OR</b>
Erythromycin.....	(base 500 mg PO QID for 7 days,	<b>OR</b>
	(ethylsuccinate) 800 mg PO QID for 7 days,	<b>OR</b>
Levofloxacin.....	500 mg PO, for 7 days,	<b>OR</b>
Ofloxacin.....	300 mg PO, BID for 7 days	

#### Syphilis (primary, secondary or early latent < 1 year)

Benzathine penicillin G.....2.4 million units IM in a single dose

#### Syphilis (latent > 1 year, latent of unknown duration, tertiary [not neurosyphilis])

Benzathine penicillin G.....2.4 million units IM for 3 doses at 1 week intervals

\*Refer to "STD Diagnostic and Treatment Guidelines" or CDC website: [www.cdc.gov/std/treatment](http://www.cdc.gov/std/treatment) for further information on treating pregnant patients, infections of the pharynx, treatment of infants and other details.